



PREVALENCE OF *ESCHERICHIA COLI* O157:H7 IN FRESH AND ROASTED BEEF IN KANO CITY, NIGERIA

***Dahiru, M., Uraih, N², Enabulele, S.A³ and Shamsudeen, U⁴.**

¹Primary Health Care Department, Dawkin Tofa Local Government Council, Kano, Nigeria.

² Department of Microbiology, Faculty of Life Sci. Univ. of Benin, Benin City.

³ Department of Biological Sciences, College of Nat. and Appl. Sci., Igbinedion Univ. Okada, Nigeria.

⁴ Department of Biological Sciences, Bayero University, Kano, Nigeria.

*Correspondence Author: Email:- musahaniya/musahaniya2@yahoo.com

ABSTRACT

The prevalence of Enterohemorrhagic *Escherichia coli* 0157:H7 in 300 fresh beef and 150 roasted beef samples from Kano city Nigeria was determined, by direct plating on Sorbitol MacConkey agar (CT-SMAC) supplemented with Cefexime 50µg/L and Potassium tellurite 25mg/L and resuscitation on Trypticase Soy Broth (TSB) at 25°C for two hours for roasted beef samples. Presumptive colonies were confirmed by using *E. coli* 0157:H7 latex agglutination test kit. Prevalence rate of 33% was obtained in fresh beef and 25.3% in roasted beef. Consumption of inadequately cooked beef poses a serious risk of infection. The study therefore stressed the need for health authorities to educate and put in place efficient surveillance strategies for detection and control of possible outbreaks of *E. coli* O157:H7 in the society.

Keywords: Prevalence, *Escherichia coli* 0157:H7, Beef

INTRODUCTION

Enterohemorrhagic *Escherichia coli* 0157:H7 is an important food-born pathogen (Benjamin and Datta, 1995) which was first identified in 1982 as a cause of hemorrhagic colitis during outbreaks of bloody diarrhea in Oregon and Michigan, USA. After these cases, several outbreaks of hemorrhagic colitis and hemolytic uremic syndrome caused by this organism have been epidemiologically linked to consumption of ground beef (Schlundt, 2001). The likely cause of *E. coli* O157:H7 infection is undercooked ground beef. Additionally, unchlorinated water, raw milk, cold sandwiches and vegetables have been implicated as sources of some outbreaks. Other implicated foods include unpasteurized apple cider and juice, salad dressing containing mayonnaise, home-made yoghurt, ozone meats, turkey roll and clams (FAO/WHO, 2003). Since then a number of works have reported the incidence of *E. coli* O157:H7 in food and during outbreaks (Hancock *et al.*, 1994; Karch *et al.* 1996; Zehn *et al.*, 2006; Agbogu *et al.*, 2006). Mead *et al.* (1999) has estimated the incidence of *E. coli* O157:H7 at 50% among EHEC serotype in relation to public health problems. This study was planned to investigate the presence of *E. coli* O157:H7 in retail fresh beef and roasted beef sold in Kano metropolis.

were randomly collected from retail markets in a sterile polyethylene bags, from the six metropolitan Local Government Areas of Kano State. The samples were analyzed within two hours of collection.

One gram (1g) of fresh beef samples each was aseptically collected placed in a sterile blender and homogenized in 9 ml of (1%) peptone water for 1 minute to obtain a homogenate. The homogenate was further serially diluted up to 10⁴ (FAO, 1979). It was then cultured on CT-SMAC for *E. coli* 0157:H7. However, for roasted beef, 1g of each sample was transferred into 9ml Trypticase Soy Broth (Lab M) shaken and incubated at 20°C for 2h (Kudo *et al.*, 2000), then 1ml each of the incubated broth was serially diluted in peptone water to 10². All the samples were cultured on Sorbitol MacConkey agar CT-SMAC (U.S Biological, Swampscott) supplemented with Cefexime (50µg/L) and Potassium tellurite (2.5 mg/L) at 45°C (March and Ratnam, 1986). The plates were allowed to solidify and incubated at 37°C for 24 hours (Chapman *et al.*, 1994; Karch *et al.*, 1996). The population of the non-sorbitol fermenters in the samples was determined by counting the colony forming unit (cfu/ml) (Wells *et al.*, 2005), and confirmed by using Latex agglutination test Kit (anti-O157:H7 antibody for *E. coli* 0157:H7) test kit DR0620M (Oxoid LTD Hampshire, England) as described by Nataro and Kaper (1998). The data obtained was analyzed by the use of ANOVA statistical analysis on SPSS package.

MATERIAL AND METHODS

Three hundred (300) samples of fresh beef and one hundred and fifty (150) roasted beef samples each

RESULTS

The result of the analysis of fresh beef and roasted beef samples is as presented in Table 1. The fresh beef has a mean NSFC count of 1.5×10^6 cfu/g and roasted beef had a mean of 3.0×10^2 cfu/g. Out of

Table 1: A comparative distribution of serologically confirmed *E. coli* 0157:H7 isolated between fresh and roasted beef

Meat sample	No. of sample Collected	Mean SFC counts (cfu/g)	Non-sorbitol fermented colonies	Serologically Confirmed	Percentages
Fresh					
Beef	300	1.5×10^6	163	158	53%
Roasted	150	3.0×10^2	38	38	25%
Beef					

Key:

SFC = Sorbitol fermenting colonies, NSFC = Non Sorbitol fermenting colonies, cfu/g = Colony forming unit per gram, EHEC = Enterohemorrhagic *E. coli* (0157:H7)

DISCUSSION

Fresh beef was found to be more contaminated than Roasted beef samples, with mean SFC counts of 1.5×10^6 cfu/g (Table 1). This might be as a result of so many factors at the slaughtering and skinning points. These include contamination from external sources like air, soil, use of non portable water and improperly washed utensils. It might also be from internal sources like intestines content, lymph nodes, as well as cross contamination by meat handlers. All these could contribute to the microbial contamination of fresh beef as described by Umoh (2001). The evidence of contamination as the most potential source of *E. coli* 0157:H7 in beef was vividly highlighted further by Elder *et al.* (2000) that meat only becomes contaminated with *E. coli* 0157:H7 when in contact with contaminated hide and or feaces during slaughter process. Evidently, *E. coli* 0157:H7 has been isolated from feaces or gastrointestinal tract of cattle, sheep, horses, pigs, turkeys, dogs and a variety of wild animals (Hancock *et al.*, 1998). However carcass contamination may vary with season, plant design and operation, geographic area, location within the plant, and to some extent, anatomical carcass site (Sofos *et al.*, 1999). The 53% prevalence rate in fresh beef is a clear indication of heavy contamination. This agrees with the work of Petridis *et al.* (2006) who reported undercooked ground beef implicated as a source of contamination that led to the infection of 243 in Montana USA. Hancock *et al.* (1994) also reported a prevalence rate of *E. coli* O157:H7 in herds of dairy cattle to be 8.3% and 16% from pastured beef cattle herds.

Although Roasted beef is least contaminated (25%) having a mean counts of 3.0×10^2 cfu/g (Table 1) and showed no significance difference between counts of samples. Uzeh *et al.* (2006) reported a total bacterial count on *Tsire-suya*, (a local delicacy) in Nigeria, to be in the range of 7×10^1 to 42×10^1 cfu/g

300 fresh beef sampled, 163 yielded NSFC out of which 158 were confirmed to be serologically positive *E. coli* 0157:H7. Of the 150 sampled roasted beef samples, 38 yielded NSFC and the 38 were confirmed to be *E. coli* 0157:H7.

an indication of high contamination, thus the mean counts of 3.0×10^2 cfu/g of NSFC shows more contamination (Uzeh *et al.*, 2006). Addition of spices is a tradition in local roasted beef preparation and have been shown to exhibit antimicrobial effects majority of which is linked to their Phenolic compounds (Shelief, 1984), however spices have played an insignificant inhibitory role, if any, as to the growth and survival of *E. coli* 0157:H7. According Frazier and Westhoff (2006) spices and condiments do not have a marked bacteriostatic activity in the concentrations normally used and unless they are treated to reduce their microbial loads, they may act high numbers and undesirable types microorganisms in the food in which they are used. Similarly, Uzeh *et al.* (2006) have reported the resistance of the activity of *Afrotomum melagueum*, *Piper guineense*, and *Capsicum frutescens* spices to *Psuedomonas aeruginosa* on *Tsire-suya*. The percentage prevalence *E. coli*/0157:H7 in roasted beef (25.3%) might be partly due to pre-enrichment (liquid and non selective medium, which enhances recovery of the injured cells (Kudo *et al.*, 2000; Clavero and Beuchat, 1995).

In addition, during roasting of beef, heat expected to have denatured microorganisms present yet *E. coli* 0157:H7 was isolated, though its optimum growth temperature is approximately 37°C (98.6°F) and is not reported to grow at temperatures below 8°C to 10°C (46°F to 50°F) or above 44°C to 45°C (Buchanan and Doyle, 1997). However, Ansary *et al.* (1999) reported *E. coli*/0157:H7 survives freezing with some decline in concentration. Equally, Clavero and Beuchat (1995), recovered 5 strains of *E. coli* 0157:H7 at 568°C for 0, 15 and 30 minutes by using non selective medium (Trypticase Soy Broth), with decrease in the microbial load in the samples, as the heating time increased. This signifies that, *E. coli* 0157:H7 could survive temperatures above 45°C.

CONCLUSION

The results of this investigation in Kano metropolis show a high prevalence rate of *E. coli* O157:H7 (53% in fresh beef and 25% in roasted beef) could be attributed to lack of good hygienic practice right from abattoir, during handling and transportation of carcass to the market. Thus, intensive sanitary measures should be taken to establish reliable hygienic standard in all operations. In view of this, health authorities and researchers have a duty to educate and put in place efficient surveillance strategies for detection and control of outbreaks of *E. coli* O157:H7 in the society.

REFERENCES

- Agbogu, V. N., Veronica, J. Umoh., Charles, A. Okuofu., Stella, I. Smith, and Joseph, B. Ameh. (2006) Study of the bacteriological and physiological indicators of pollution of surface waters in Zaria, Nigeria. *Afr. J. of Biotech.*, Vol. **5** (9): 732-737
- Agaoglu ,S., Yavuz, M.T., Berktaş, M. and Gündüçoglu, H. (2000) Defection of *Escherichia coli* in Retail Ground Beef, Raw Ground Beef, Patties and Raw Meat , Balls sold in van. *East. J. of Med.* **5**(2): 7375.
- Ansay, S.E., Darling, K.A. and Kaspar, C.W. (1999) Survival of *Escherichia coli* O157:H7 in ground-beef patties during storage at 2, -2, 15 and then -2 degrees C, and -20 degrees. *C. J. Food Prot* **62**(11):1243-7.
- Benjamin, M. M. and A. R. Datta. (1995). Acid tolerance of enterohemorrhagic *Escherichia coli*. *Appl. Environ. Microbiol.* **61**:1669-1672
- Buchanan, R. L. and M. P. Doyle. (1997). *Food borne Disease Significance of Escherichia coli O157:H7 and other Enterohemorrhagic E. coli*. *Food Technology* **51**: 69-75.
- Chapman, P. A., D. J. Wright, and C. A. Siddons. (1994). A comparison of immunomagnetic separation and direct culture for the isolation of verocytotoxin-producing *Escherichia coli* O157 from bovine faeces. *J. Med. Microbiol.* **40**:424-427.
- Clavero, M. R. S. and L. R. Beuchat. (1995). Suitability of Selective Plating Media for recovering Heat- or Freeze-Stressed *Escherichia coli* O157:H7 from Tryptic Soy Broth and Ground Beef; *J. Appl. and E. Microb.* **61** (9): 3268-3273
- Elder, R. O., J. E. Keen., G. R. Siragusa., G. A. Barkocy-Gallagher., M. Koohmaraie, and W. W. Laegreid. (2000). Correlation of enterohemorrhagic *Escherichia coli* O157 prevalence in feces, hides, and carcasses of beef cattle during processing. *Proc. Natl. Acad. Sci.* **97**:2999-3003.
- Food and Agricultural Organization (FAO), (1979). *Manuals of Food quality Control for Microbiological Analysis*.
- Food and Agricultural Organization/World Health Organization (FAO/WHO), Codex Committee on Food Hygiene, (2003). Risk profile for Enterohemorrhagic *Escherichia coli*, including the identification of the commoditis of concerned, including sprouts, ground beef and pork. CX/FH 03/5-Add.4 Sep. 2002.
- Frazier, W.C. and Westhoff, S.A.(2006) *Food Microbiology* 3rd Edition McGraw Hill Publishing Company Limited. Newyork, pp.163-165,223-236,419-543
- Hancock, D. D., T. E. Besser., M. L. Kinsel., P. I. Tarr., D. H. Rice, and M. G. Paros. (1994). The prevalence of *Escherichia coli* O157:H7 in dairy and beef cattle in Washington state. *Epidemiol. Infect.* **113**:199-207.
- Hancock, D. D., Besser, T. E. and Rice, D.G. (1998). Multiple source of *Escherichia coli* O157 in feed lots and dairy farms in the northwestern USA. *Prev. Vet. Med.* **35**: 11-19.
- Kardch, H., C. Janetzki-Mittmann., S. Aleksic. and M. Datz. (1996). Isolation of enterohemorrhagic *Escherichia coli* O157 strains from patients with hemolytic-uremic syndrome by using immunomagnetic separation, DNA-based methods, and direct culture. *J. Clin. Microbiol.* **34**:516-519
- Kudo, H. Y., M. Ikeda., H. Kodaka., H. Nakagawa., K. Goto., T. Masuda., H. Konuma., T. Kojima. and S. Kumagai. (2000) Selective Enrichment with a Resuscitation Step for Isolation of Freeze-Injured *Escherichia coli* O157:H7 from Foods. *J. Appl. and Environ. Microb.* **66** (7): 2866-2872.
- Kudva, I. T., P. G. Hatfield, and C. J. Hovde. (1995). Effect of diet on the shedding of *Escherichia coli* O157:H7 in a sheep model. *Appl. Environ. Microbiol.* **61**:1363-1370. March
- March, S.B. and Ratnam, S. (1986) Sorbitol-MacConkey medium for detection of *Escherichia coli* O157:H7 associated with hemorrhagic colitis. *J. Clin Microbiol.* **23**:869-872.
- Mead, P.S., Slutsker, L., Dietz, V., L.F. McCraig., J.S Bressee., C. Shapiro., P. M. Griffin., and Tauxe. (1999). Food related illness and death in the united states. *J. Emerg. Infect. Dis.* **5**(5): 607-25.
- Nataro, J. P., and J. B. Kaper. (1998). Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.* **11**:142-201.
- Petridis, H., G. Kidder, and A. Ogram. (2006). *Escherichia coli* O157:H7 A Potential Health Concern. University of Florida IFAS Extension.
- Schlundt, J. (2001). Emerging food borne pathogens. *Biomed. J. Environ. Sci.* **14**(1-2):44-52.
- Shelef, L.A. (1984). Antimicrobial effects of spices. *J. of Food Safety*. **6**(1) 29-44

- Shin Sata, Tomohiko, Fujisawa,, Ro Osawa,, Atsushi Iguchi., Shirio Yamai, and Toshio, Shimada. (2003). An Improved Enrichment Broth for Isolation of *Escherichia coli* O157, with Specific Reference to Starved Cells, from Radish Sprouts. *Applied and Environ. Micro.* Vol. **69**, No. 3. 1858–1860
- Sofos, J. N., Kochavar, S. L., Bellinger, G. R., Buege, D. R., Hancock, D. D., Ingham, S. C., Morgan, J. B., Reagan, J. O. and Smith, G. C. (1999) Sources and extent of microbiological contamination of beef carcasses in seven United States slaughtering plants. *J. Food Prot.* **62**:140-145.

- Umoh, J.U. (2001). An overview of possible control points of ready-to-eat beef products in northern Nigeria. International conference on food and security, conference center Ib Nigeria. Pp 109-115
- Uzeh, R.E., R.E.O. henhen, and O.O.A dentii. (2002) Bacterial contamination of 'Tsire- Suya' Nigerian meat product. *Pakistan J. Nutrition* **5**(5):458-460
- Wells, J. E., Barry, E.D. and Varel, W.H. (2005) Effect of common forage phenolic acids on *Escherichia coli* O157:H7 viability in cattle feces. *J.Appl.Environ. Microbiol.* **71**(12):7974-7979.